

Peierls Dimerization with Non-Adiabatic Spin-Phonon Coupling

G. Wellein¹, H. Fehske¹, and A.P. Kampf²

¹ *Physikalisches Institut, Universität Bayreuth, 95440 Bayreuth, Germany*

² *Institut für Physik, Theoretische Physik III, Elektronische Korrelationen und Magnetismus, Universität Augsburg, 86135 Augsburg, Germany*

We study the magnetic properties of a frustrated Heisenberg spin chain with a dynamic spin-phonon interaction. By Lanczos diagonalization, preserving the full lattice dynamics, we explore the non-adiabatic regime with phonon frequencies comparable to the exchange coupling energy which is e.g. the relevant limit for the spin-Peierls compound CuGeO₃. When compared to the static limit of an alternating spin chain the magnetic properties are strongly renormalized due to the coupled dynamics of spin and lattice degrees of freedom. The magnitude of the spin triplet excitation gap changes from a strong to a weak dimerization dependence with increasing phonon frequencies implying the necessity to include dynamic effects in an attempt for a quantitative description of the spin-Peierls state in CuGeO₃.

PACS numbers: 75.10.Jm, 75.40.Mg, 75.50.Ee

Low dimensional quantum spin systems have attracted considerable attention of theorists over the decades. Most of the remarkable features observed in these systems are pure quantum effects uniquely due to their low dimensionality. Among the current experimental and theoretical efforts for understanding the magnetic properties of linear chain and ladder materials the recent discovery of a spin-Peierls (SP) transition at $T_{SP} = 14.3K$ in the inorganic compound CuGeO₃ (CGO) [1] has received particular attention. In analogy to the Peierls instability towards dimerization in quasi one-dimensional metals [2] the energy of spin chains is lowered by dimerizing into an alternating pattern of weak and strong bonds. The SP transition is therefore driven by the magnetic energy gain which overcompensates the lattice deformation energy [3]. In the SP phase of CGO the copper moments form singlet dimers along the chains and there is an energy gap to spin triplet excitations. Experimentally, the SP nature of the transition and the spin gap have been firmly established by inelastic neutron scattering (INS), susceptibility, X-ray, and electron-diffraction experiments [4].

The Peierls transition naturally involves also the lattice degrees of freedom due to the alternating distortion of atomic positions. A rather common situation has been the clear separation of the electronic or magnetic energy scales from the frequencies of those phonons which couple most effectively to the SP lattice distortion. This adiabatic limit is e.g. realized in trans-polyacetylene [5] or organic SP materials [6]. In contrast, in CGO two weakly dispersive optical phonon modes involved in the SP transition have been identified with frequencies $\hbar\omega \approx J$ and $\hbar\omega \approx 2J$, where J is the superexchange coupling between neighboring Cu spins [7]. These hard phonons require a very strong spin-phonon coupling for a SP transition to occur [8]. A strong magnetoelastic coupling has indeed been observed in thermal expansion and magnetostriction experiments [9] and inferred from large uniaxial pressure (p_i) derivatives $\partial J/\partial p_i$ [10]. The SP

physics in CGO is therefore in a non-adiabatic regime.

Despite this unusual situation previous theoretical studies have commonly adopted an alternating and frustrated antiferromagnetic (AF) Heisenberg spin chain model [11]

$$H = J \sum_i [(1 + \delta(-1)^i) \mathbf{S}_i \cdot \mathbf{S}_{i+1} + \alpha \mathbf{S}_i \cdot \mathbf{S}_{i+2}] \quad (1)$$

with a static dimerization parameter δ , thus representing the extreme adiabatic limit of a SP chain. i denotes the sites of a chain with length N and $\mathbf{S}_i$ are $S = 1/2$ spin operators. α determines the strength of a frustrating AF next-nearest neighbor coupling. The model (1) contains two independent mechanisms for spin gap formation. At $\delta = 0$ and for $\alpha < \alpha_c$ the groundstate is a spin liquid and the elementary excitations are massless spinons [12]. $\alpha_c = 0.241$ was accurately determined by numerical studies [13,11]. For $\alpha > \alpha_c$ the groundstate is spontaneously dimerized, the spectrum acquires a gap and the elementary excitations are massive spinons [14,15]. On the other hand for any finite δ the singlet groundstate of (1) is also dimerized, but the elementary excitation is a massive magnon [12,16].

From fits to magnetic susceptibility data in the uniform high temperature phase [17,10] and from the requirement to reproduce the experimental triplet excitation gap [17,18] the parameter set $J = 160K$, $\alpha = 0.36$, and $\delta = 0.014$ was estimated for CGO within the static model (1). Yet, the amplitude of the dimerization is substantially underestimated when compared to estimates from structural data in the SP phase [19,7]. In this letter we will show that due to the non-adiabatic phonon dynamics a much larger dimerization is needed to achieve the same magnitude of the triplet gap and thus offers a possible route towards a quantitatively consistent description.

The dimerization parameter δ in Eq. (1) may be viewed to result from a mean-field treatment of an interchain




FIG. 1. Energies of the singlet groundstate (full symbols) and the lowest triplet excitation (open symbols) vs. the number of phonons used in the numerical diagonalization. The different symbols indicate the results obtained by including phonons (a) only with $q = \pi$ ($\circ$), (b) with $q = 0$ and $q = \pi$ ($\diamond$), and (c) at all wavevectors ($\square$). As in all subsequent figures energies are measured in units of J .

coupling which is generated by a purely elastic mechanism [20]. In fact, this coupling to neighboring dimerized chains provides an external potential lifting the degeneracy between the two possible dimerized groundstates [21]. Thereby, soliton and antisoliton excitations form spin triplet or spin singlet bound states [22], the former is the elementary massive magnon of the spin Hamiltonian (1) for $\delta > 0$.

In this letter we choose a different starting point and use a Heisenberg chain model in which the spins are explicitly coupled to the lattice degrees of freedom maintaining their full quantum dynamics. Specifically we study the magnetic properties of the Hamiltonian [21]

$$H_{SP} = J \sum_i [(1 + g(b_i^+ + b_i)) \mathbf{S}_i \cdot \mathbf{S}_{i+1} + \alpha \mathbf{S}_i \cdot \mathbf{S}_{i+2}] + \hbar\omega \sum_i b_i^+ b_i \quad (2)$$

with the local phonon creation and annihilation operators b_i^+ and b_i , respectively. For the optical phonon with frequency ω we thus choose a dispersionless Einstein mode – an appropriate choice for modeling those phonons in CGO which are most relevant for the Peierls distortion [7]. In the non-adiabatic ($\hbar\omega \sim J$) and intermediate coupling ($g \sim J$) regime the spin and the phonon dynamics are intimately coupled and integrating out the phonons in order to obtain an effective renormalized spin-only Hamiltonian is in general not possible [23].

This situation thus requires a numerical approach. We have performed exact diagonalizations on finite chains up to $N = 16$ sites with periodic boundary conditions.




FIG. 2. Finite size scaling behavior of the triplet excitation gap Δ^{ST} for $\alpha = 0.36$. The lines result from the fit to the scaling function (3). The (almost adiabatic) phonon parameters $g = 0.11$ and $\omega = 0.3$ were used in order to make contact with previous numerical studies [25].

Since the Hilbert space associated with the phonons is infinite even for a finite system, we apply a controlled truncation procedure retaining only basis states with at most M phonons [24]. M is fixed by requiring a relative error in the groundstate energy less than 10^{-7} . In our calculations all possible phonon modes are taken into account.

Fig. 1 demonstrates for a 12 sites chain how many phonons have to be included to achieve convergence for the groundstate energies of the singlet $S = 0$ and the triplet $S = 1$ sectors. Although the specific value of α is not crucial for our arguments, we keep the frustration parameter $\alpha = 0.36$ in all calculations fixed to its most likely value in CGO. Fig. 1 contrasts the results obtained if only phonons with wavevector $q = \pi$ or $q = \pi$ and $q = 0$ are used to the case when all phonon modes are included. While it has been argued in Ref. [25] that keeping the $q = \pi$ mode only should already capture the dominant dimerization effect of the spin-phonon interaction, we find in Fig. 1 that e.g. the singlet-triplet excitation gap $\Delta^{ST} = E_0^T - E_0^S$ is strongly renormalized when phonons of *all* q are taken into account implying that a spin triplet excitation is accompanied by a *local* distortion in the lattice.

The renormalization of the singlet-triplet gap Δ^{ST} is shown in the scaling plot in Fig. 2. It is contrasted to the result with a restriction to the $q = \pi$ phonons only. Despite of the limited system sizes of up to 16 sites the data are well fitted by the scaling function [27,18]

$$\Delta^{ST}(N) = \Delta^{ST} + \frac{A}{N} e^{-N/N_0}. \quad (3)$$

The data clearly demonstrate that the restriction to the π -modes substantially overestimates the size of the gap




FIG. 3. Dispersion of the elementary triplet excitation (left scale) for a chain with $N = 16$ sites for different phonon frequencies and spin-phonon coupling constants. Experimental data (*, right scale) are also included from Ref. [26].

for $N \rightarrow \infty$.

The triplet dispersion shown in Fig. 3 supports the expectation – as inferred from the neutron scattering studies of the phonons [7] – that the frequencies of the relevant optical phonon modes in CGO are comparable to or larger than the exchange coupling J . Because for $\hbar\omega < J$ the phononic character of the excitation dominates leading to a flattening of the dispersion [28,29]. This, however, is incompatible with the magnetic INS data [26] which do not show any signature for a flat dispersion of the triplet excitation branch. We have included in Fig. 3 experimental data on CGO from Ref. [26] which, however, is not intended as a fit to the data – our results in Fig. 3 are for a fixed chain length with $N = 16$ sites – but rather shows the shape of the triplet dispersion. We stress that the upturn at $Q = 0$ is a finite-size effect which also occurs for the pure spin model and rapidly decreases with increasing N [30]; for $N \rightarrow \infty$, $E_0^T(\pi)$ and $E_0^T(0)$ become degenerate.

The Peierls ordering structure of the groundstate can be demonstrated in two complementary ways. The *magnetic* ordering pattern of an alternating local singlet formation is captured by the order parameter

$$D = \langle \frac{1}{N} \sum_i (-1)^i (\mathbf{S}_{i-1} \cdot \mathbf{S}_i - \mathbf{S}_i \cdot \mathbf{S}_{i+1}) \rangle. \quad (4)$$

D is the proper choice for the Heisenberg spin chain model (1). For the translationally invariant quantum spin-phonon Hamiltonian (2) the Peierls distortion of the lattice is reflected in the spatial structure of the displacement correlation function

$$C_{1,i} = \langle (b_1 + b_1^\dagger) (b_i + b_i^\dagger) \rangle - \delta_{1,i}. \quad (5)$$

The alternating structure of the correlation function $C_{1,i}$ shown in Fig. 4 implies the Peierls formation of short




FIG. 4. Displacement correlation function $C_{1,i}$ as defined in the text for different phonon frequencies and spin lattice coupling constants for a chain with $N = 12$ sites.

and long bonds and thus alternating strong and weak AF exchange interactions, i.e. a dimerized groundstate. Naturally the dimerization is enhanced (weakened) by increasing the spin-phonon coupling (phonon frequency). We note that for any finite phonon frequency there exists a critical coupling constant g_c [31]; in the infinite system quantum lattice fluctuations will destroy the Peierls dimerization for couplings $g < g_c$. For non-adiabatic phonon frequencies $\hbar\omega \sim J$ also the critical coupling g_c is comparable to J – a situation which matches the strong magnetoelastic coupling and the SP physics in CGO.

In order to quantify these results we obtain the magnitude of the Peierls dimerization order parameter δ from the static (lattice) structure factor at wavenumber $q = \pi$

$$\delta^2 = \frac{g^2}{N^2} \sum_{i,j} \langle u_i u_j \rangle e^{i\pi(R_i - R_j)} \quad (6)$$

where $u_i = b_i + b_i^\dagger$. Like in the ordinary Peierls phenomenon, a finite dimerization $\delta > 0$ necessarily leads to a gap Δ^{ST} in the magnetic excitation spectrum. For evaluating the relation between the dimerization and the resulting magnitude of the spin triplet excitation gap we keep the phonon frequency fixed, vary the coupling strength g and calculate for each parameter set Δ^{ST} (see Fig. 5), and the dimerization δ from Eq. (6).

The results for the static (spin-only) and the dynamic spin-phonon model are compared in Fig. 5 for three different chain lengths. For vanishing lattice dimerization, i.e. in the absence of any spin-phonon coupling ($g = 0$), the results for the static and the dynamic model naturally agree (note that for $\delta = 0$ there remains a spin excitation gap even for $N \rightarrow \infty$ due to the frustration driven singlet dimer ordering $D \neq 0$). The important message of Fig. 5 is that a substantial enhancement of the dimerization is needed to achieve a given triplet excitation gap, if




FIG. 5. Triplet excitation gap Δ^{ST} versus dimerization δ for the static spin-only ($\omega = 0$, $\diamond \square \circ$) and the dynamic spin-phonon model (full symbols) with phonon frequency $\omega = 1$, for different chain lengths at $\alpha = 0.36$. The inset shows the results for the dynamic eight-site model (filled symbols; $\omega = 0.1$ [0.3] triangles up [down], 1.0 ($\bullet$)) in comparison to the static case ($\circ$).

the dynamic phonons have frequencies comparable to J . This partially resolves the quantitative problem encountered within the alternating and frustrated Heisenberg spin chain model (1), because fixing parameters J and α by fitting the high temperature susceptibility and δ by matching Δ^{ST} to the INS data in CGO, the dimerization parameter δ is underestimated by roughly a factor of 5 [19,32]. The inset in Fig. 5 shows the evolution of the Δ^{ST} vs. δ results with increasing phonon frequency ω at a fixed chain length. With decreasing ω the data smoothly merge with the result for the static spin-only model.

In conclusion, we have numerically analyzed the Peierls dimerized groundstate and the spin triplet excitations in a Heisenberg chain model with dynamic spin-phonon coupling. The magnetic excitations inherently include a local lattice distortion requiring a multi-phonon mode treatment of the lattice degrees of freedom. A spin-phonon coupling induced flattening of the triplet dispersion is avoided for a strong spin-phonon coupling in the non-adiabatic regime. The non-adiabatic phonon dynamics strongly renormalizes the magnetic excitation spectrum and the dimerization dependence of the triplet excitation gap. These features are in accordance with experimental data on CGO suggesting the necessity for a non-adiabatic spin-phonon approach to the SP physics in this material.

We acknowledge useful discussions with G. Bouzerar, M. Braden, D. Khomskii, W. Weber, and A. Weiße. We are particularly indebted to B. Büchner for informing us about unpublished estimates for the dimerization of J in the SP phase of CGO. The numerical calculations were

performed on the vector-parallel supercomputer Fujitsu VPP700 at the LRZ München.

-
- [1] M. Hase et al., Phys. Rev. Lett. **70**, 3651 (1993).
 - [2] R.E. Peierls, *Quantum Theory of Solids*, Clarendon, Oxford, 1955.
 - [3] M.C. Cross and D.S. Fisher, Phys. Rev. B **19**, 402 (1979).
 - [4] For a review see J.P. Boucher and L.P. Regnault, J. Phys. I (France) **6**, 1 (1996).
 - [5] A.J. Heeger et al., Rev. Mod. Phys. **60**, 781 (1988).
 - [6] For a review see J.W. Bray et al., in *Extended Linear Chain Compounds*, J.S. Miller (Ed.), Plenum, New York, 1983, Vol. 3, pp. 353–415.
 - [7] M. Braden et al., Phys. Rev. B **54**, 1105 (1996); and to be published in Phys. Rev. Lett..
 - [8] L. Bulaevskii et al., Sol. State Comm. **13**, 125 (1973).
 - [9] B. Büchner et al., Phys. Rev. Lett. **77**, 1624 (1996).
 - [10] K. Fabricius et al., Phys. Rev. B **57**, 1102 (1998).
 - [11] G. Castilla et al., Phys. Rev. Lett. **75**, 1823 (1995).
 - [12] F.D.M. Haldane, Phys. Rev. B **25**, 4925 (1982).
 - [13] K. Okamoto and K. Nomura, Phys. Lett. A **169**, 433 (1993).
 - [14] R. Chitra et al., Phys. Rev. B **52**, 6581 (1995).
 - [15] S.R. White and I. Affleck, Phys. Rev. B **54**, 9862 (1996).
 - [16] A. M. Tsvelik, Phys. Rev. B **45**, 486 (1992).
 - [17] J. Riera and A. Dobry, Phys. Rev. B **51**, 16098 (1995).
 - [18] G. Bouzerar et al., preprint cond-mat/9701176.
 - [19] B. Büchner, private communication: From the uniaxial pressure derivatives of the exchange coupling J [10] and the structural distortion in the dimerized phase [7] it is possible to show that the dimerization of J is significantly stronger than $\delta = 1.4\%$. The result of a straightforward data analysis leads to an estimate $\delta \approx 4\%$ – 5% .
 - [20] F.H.L. Essler et al., Phys. Rev. B **56**, 11001 (1997).
 - [21] D. Khomskii et al., Czech. Journ. of Phys. **46**, Suppl. S6, 32 (1996).
 - [22] G. Bouzerar et al., cond-mat/980146.
 - [23] An attempt in this direction has recently been formulated by G. Uhrig, preprint cond-mat/9801185.
 - [24] B. Bäuml et al., preprint cond-mat/9802181.
 - [25] D. Augier and D. Poilblanc, Eur. Phys. Jour. B **1**, 19 (1998).
 - [26] L.P. Regnault et al., Phys. Rev. B **53**, 5579 (1996).
 - [27] T. Barnes and J. Riera, Phys. Rev. B **50**, 6817 (1994).
 - [28] We emphasize that the flattening of the dispersion can not be obtained if only phonon modes with $q = \pi$ are included as in Ref. [25].
 - [29] G. Wellein and H. Fehske, Phys. Rev. B **56**, 4513 (1997).
 - [30] G. Wellein, PhD thesis, Universität Bayreuth, 1998.
 - [31] D. Augier et al., preprint cond-mat/9802053.
 - [32] M. Braden, private communication.